


Process for the Preparation of Fuopyrroles

The present invention relates to a microwave assisted rapid and economical process for the preparation of fuopyrroles of the general formula I, comprising (a) heating a compound of the formula II under microwave irradiation optionally in the presence of an inert solvent. The fuopyrroles of the general formula I can be obtained in high yield and high purity by the process of the present invention.


WO03022848 discloses a process for the preparation of fuopyrroles of the general formula I, comprising heating a compound of the formula




wherein A¹ and A² have the meanings as given below and R is C₁-C₁₈alkyl, in particular C₁-C₄alkyl, aryl, in particular phenyl, or aralkyl, in particular benzyl, which can be substituted one to three times with C₁-C₈alkyl, C₁-C₈alkoxy, or halogen. Examples of inert solvents include, but are not limited to, aromatic solvents, like biphenyl, para-, meta or ortho-terphenyl, dibenzyltoluene, α-methyl- or β-methylnaphthalene, cyclic carbonates, like 1,3-dioxolan-2-one, ketones, like acetophenone or benzophenone, γ-butyrolactone and ethylene glycols, like Phe-Cellosolve or Bu-Cellosolve, or mixtures thereof, in particular mixtures of di- and triarylethers (Dowtherm A®).

It has now surprisingly been found, that the 3,6-diphenylfuro[3,4-c]pyrrole-1,4-diones (fuopyrroles) of formula I can be obtained in higher yield by carrying out the above reaction under microwave radiation. The yield of the ring closure of ethyl 4-benzoyl-4,5-dihydro-5-oxo-2-phenylpyrrole-3-carboxylate to 3,6-diphenylfuro[3,4-c]pyrrole-1,4-dione is, for example, increased from 40 to 86 % by the microwave assisted process according to the present invention. Moreover, we have observed that the preparation of this lactone (a versatile DPP precursor) requires lesser time (1 to 10 minutes) under microwave irradiation while ring closure of the compound of formula II takes 60 hours when conducted without microwave radiation (conventional method). In addition, the solvent can be omitted in the microwave assisted ring closure, which makes the above process further cost effective.

Accordingly, the present invention relates to a process for the preparation of furopyrroles of



(a) heating a compound of the formula



5 solvent,

wherein A¹ and A² are C₁-C₁₈alkyl, C₂-C₁₈alkenyl, C₂-C₁₈alkynyl, C₅-C₈cycloalkyl, C₅-C₈cycloalkenyl, aryl or heteroaryl,

A³ is hydrogen, C₁-C₁₈alkyl, cyanomethyl, Ar³, -CR³⁰R³¹-(CH₂)_m-Ar³ or Y-R³², wherein R³⁰ and R³¹ independently of each other stand for hydrogen or C₁-C₄alkyl, or phenyl which can be

10 substituted up to three times with C₁-C₄alkyl,

Ar³ stands for aryl, C₅-C₈cycloalkyl, C₅-C₈cycloalkenyl or heteroaryl, which can be substituted one to three times with C₁-C₈alkyl, C₁-C₈alkoxy, halogen or phenyl, which can be substituted with C₁-C₈alkyl or C₁-C₈alkoxy one to three times, and m stands for 0, 1, 2, 3 or 4,

R is C₁-C₁₈alkyl, in particular C₁-C₄alkyl, aryl, in particular phenyl, or aralkyl, in particular

15 benzyl, which can be substituted one to three times with C₁-C₈alkyl, C₁-C₈alkoxy, or halogen, Y is -C(O)-, -C(O)O-, -C(O)NH-, -SO₂NH- or -SO₂- and

R³² is C₁-C₁₈alkyl, Ar³, or aralkyl.

If desired, the process of the present invention can be carried out in the presence of an inert solvent. Examples of inert solvents include, but are not limited to, aromatic solvents, like

20 biphenyl, para-, meta or ortho-terphenyl, dibenzyltoluene, α-methyl- or β-methylnaphthalene, cyclic carbonates, like 1,3-dioxolan-2-one, ketones, like acetophenone or benzophenone, γ-butyrolactone and ethylene glycols, like Phe-Cellosolve or Bu-Cellosolve, or mixtures thereof, in particular mixtures of di- and triarylethers (Dowtherm A®).

In a preferred embodiment the compound of the formula II is heated for about 1 to 60

25 minutes at a temperature of 180 to 280 °C, preferably 180-230 °C, with or without solvent, under microwave irradiation.


A microwave furnace suitable for the irradiating the composition comprises a microwave source, microwave frequency range selector, a microwave frequency modulator to modulate the microwave frequency across the selected frequency range, microwave forward power controller to select the forward power setting, a thermocouple, an infrared temperature sensor or other temperature measuring means, and a microwave forward power on/off controller to turn the forward power on and off in response to the temperature of the composition. Frequency modulation increases the uniformity of the power distribution throughout the furnace cavity, thereby heating the composition uniformly. Suitable microwave furnaces are described in, for example, U.S. Pat. Nos. 5,321,222 and 5,961,871 to Bible et al., U.S. Pat. No. 5,648,038 to Fathi et al., and U.S. Pat. No. 5,521,360 to Johnson et al. A presently preferred microwave furnace is commercially available from CEM, Inc., as model Discover®. The Discover® System incorporates temperature and pressure feedback systems, for example, an infrared temperature sensor positioned below the reaction vessel, for complete control of the reaction.

It is preferred that the reaction mixture be irradiated in a vessel transparent to microwave radiation in the frequency range employed.

The samples, comprising the compounds of formula II and optionally the solvent, are advantageously heated in pressurized tubes, such as, for example, sealed glass tubes, whereby the pressure is allowed to increase up to $25 \cdot 10^5$ Pa. Preferably the pressure is between 1 to $14 \cdot 10^5$ Pa.

The selection of the actual microwave frequency range will depend on the reactants, but will generally be about 0.9 to about 2.45 GHz. Selection of a forward power input will depend on the nature of the reactants. For example, in the synthesis of 3-(*p*-bromophenyl)-6-phenyl furo[3,4-*c*]pyrrole-1,4-dione, a preferred forward power level is about 150 to 300 watts.

As described in WO03022848 the furopyrroles of formula I can be used as crystal growth regulators and are intermediates in the synthesis of diketopyrrolopyrroles, which can be obtained by reacting a compound of formula I with a primary amine of the formula A^4-NH_2



wherein A^4 is C_1-C_{18} alkyl or Ar^3 , and A^1 , A^2 and A^3 are as defined above.


The reaction between the compound of the general formula I and the primary amine or the mixture of primary amines is carried out in a suitable inert solvent or dispersant.

- Suitable solvents or dispersants are, for example, ethers, in particular those having 2 to 8 carbon atoms in the molecule, such as, for example, diethyl ether, methyl ethyl ether, di-n-propyl ether, diisopropyl ether, methyl n-butyl ether, methyl tert-butyl ether, ethyl n-propyl ether, di-n-butyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane, bis-β-methoxyethyl ether; oligoethylene glycol dimethyl ethers, such as, for example, pentaglyme; aliphatic hydrocarbons, such as, for example, hexane, heptane, low- and high-boiling petroleum ethers; cycloaliphatic hydrocarbons, such as, for example, cyclohexane, methylcyclohexane, tetralin, decalin; aromatic hydrocarbons, such as, for example, benzene, toluene, o-, m- and p-xylene, ethylbenzene; halogenated aliphatic or aromatic hydrocarbons, such as, for example, methylene chloride, chloroform, carbon tetrachloride, chlorobenzene, dichlorobenzene; nitriles, such as, for example, acetonitrile; amides, such as, for example, dimethylformamide, dimethylacetamide, N-methylpyrrolidone; hexamethylphosphoric triamide; and sulfoxides, such as, for example, dimethyl sulfoxide. Mixtures of various solvents can also be used.
- The reaction is preferably carried out in a dipolar or non-polar aprotic solvent. Examples of preferred aprotic solvents are: dimethylformamide, dimethyl sulfoxide, hexamethylphosphoric triamide, sulfolane, N-methylpyrrolidone, tetramethylurea, acetonitrile, ethylene glycol dimethyl ether, ethylene glycol diethyl ether, diethylene glycol dimethyl ether and triethylene glycol dimethyl ether, nitromethane, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU), 1,3-dimethyl-2-imidazolidinone, benzonitrile, nitrobenzene, chloroform, carbon tetrachloride and methylene chloride. Particularly preferred aprotic solvents are chloroform, carbon tetrachloride and methylene chloride, of which chloroform is particularly preferred. The reaction between the compound of the general formula I and the primary amine IV is carried out in the presence of a dehydrating agent. Examples of suitable dehydrating or water-eliminating agents of this type are: N,N'-disubstituted carbodiimides, in particular if they contain at least one secondary or tertiary alkyl radical, such as, for example, diisopropyl-, dicyclohexyl- or N-methyl-N'-tert.-butylcarbodiimide (cf. "The Chemistry of Ketenes, Allenes and Related Compounds", Part 2, Editor: S. Patai, John Wiley & Sons 1980, 722-753). Dicyclohexylcarbodiimide is particularly suitable.
- The reaction between the compound of the formula I and the primary amine IV can be carried out, for example, at temperatures from -10° C up to the boiling point of the solvent or solvent mixture used. In many cases it is carried out at -10 to 30 °C and preferably at room temperature. 0.9 to 1.4 mol, preferably 1.0 to 1.3 mol of the primary amine IV are in general employed per mole of compound of the general formula I. The reaction can be catalyzed by adding a strong non-aqueous acid such as trifluoroacetic acid.

5

The primary amines IV are known or can be easily prepared by the methods known for the preparation of these class of compound.


The starting compound of the formula Ia, wherein A³ is different from a hydrogen atom, is obtained by reacting a compound of the formula



A³ have the meanings as given above and X is a leaving group. The reaction between the compound of the general formula Ia and the compound of the formula V is carried out in a suitable inert solvent such as tetrahydrofuran, in the presence of a base such as sodium hydride (NaH) or sodium hexamethyldisilazane (NaHMDS), at a temperature ranging from 20 °C to the boiling point of the solvent. The term "leaving group" means a group, such as iodo, bromo or chloro, benzene- or p-toluenesulfonate. Processes for the introduction of A³ into compounds of the formula Ia are described, for example, in US-A-4,585,878.

Suitable alkylating agents are, for example, alkyl halides, in particular alkyl iodides, reactive alkyl esters, in particular alkyl esters of sulfonic acids, such as, for example, alkyl esters of benzene- or p-toluenesulfonic acid. Suitable arylating agents are for example activated aryl compounds such as 1-fluoro-2,4-dinitro-benzene.

The starting compound of the formula IIa is obtained by reacting a compound of the formula




base, such as for example NaH or NaHMDS at a temperature ranging from 25 °C to the boiling point of the solvent, wherein R, A¹ and A² have the meanings as given above.

The starting compounds of the formula VI are known or can be prepared in analogy to processes described in US-A-4,681,971, US-A-4,749,795, US-A-4,720,305 and US-A-4,659,775.


Alternatively, compounds of the formula

6




(VIII), wherein A³ is different from a hydrogen atom and is in particular


aryl, can be prepared by a copper catalyzed decomposition of diazoacetates in the presence of enaminoamides (G. Maas, A. Müller, J. prakt. Chem. 340 (1998) 315-322):



- 5 In addition, compounds of formula (VIII) wherein A³ is aryl can be obtained by reacting a compound of formula (IIb) with an amine A³-NH₂:



- 10 Preferably, the lactone of formula (IIb) is reacted with aniline to afford the N-phenyl pyrrolinone ester of formula (VIII) as described in more detail in Example 4. The compounds of the formula VI, wherein A³ is different from a hydrogen atom and is in particular aryl, can be reacted to compounds of the formula III as described above.



In addition, DPP of formula (III) wherein A¹ and A² are C₁-C₁₈alkyl, C₂-C₁₈alkenyl, C₂-C₁₈alkynyl, C₅-C₈cycloalkyl, C₅-C₈cycloalkenyl, aryl or heteroaryl,

A³ is hydrogen, C₁-C₁₈alkyl, cyanomethyl, Ar³, -CR³⁰R³¹-(CH₂)_m-Ar³ or Y-R³², wherein R³⁰ and R³¹ independently of each other stand for hydrogen or C₁-C₄alkyl, or phenyl which can be substituted up to three times with C₁-C₄alkyl,


Ar³ stands for aryl, C₅-C₈cycloalkyl, C₅-C₈cycloalkenyl or heteroaryl, which can be substituted one to three times with C₁-C₈alkyl, C₁-C₈alkoxy, halogen or phenyl, which can be substituted with C₁-C₈alkyl or C₁-C₈alkoxy one to three times, and m stands for 0, 1, 2, 3 or 4,

R is C₁-C₁₈alkyl, in particular C₁-C₄alkyl, aryl, in particular phenyl, or aralkyl, in particular benzyl, which can be substituted one to three times with C₁-C₈alkyl, C₁-C₈alkoxy, or halogen, Y is -C(O)-, -C(O)O-, -C(O)NH-, -SO₂NH- or -SO₂-,

R³² is C₁-C₁₈alkyl, Ar³, or aralkyl, and

A⁴ is hydrogen

can also directly be obtained by reacting a compound of formula (VIII) with a nitrile A²-CN, wherein A¹, A² and A³ have the meanings as given above:



Further, the compound of the formula III can be reacted with a compound of the formula A^5-X , wherein A^5 has the meaning of A^3 as given above and X is a leaving group. The reaction between these compounds is carried out in a suitable inert solvent such as tetrahydrofuran, in the presence of a base such as sodium hydride (NaH) or sodium


5 hexamethyldisilazane (NaHMDS), at a temperature ranging from 20 °C to the boiling point of the solvent. The term "leaving group" means a group, such as iodo, bromo or chloro, benzene- or p-toluenesulfonate.

Suitable alkylating agents are, for example, alkyl halides, in particular alkyl iodides, reactive
10 alkyl esters, in particular alkyl esters of sulfonic acids, such as, for example, alkyl esters of benzene- or p-toluenesulfonic acid. Suitable arylating agents are for example activated aryl compounds such as 1-fluoro-2,4-dinitro-benzene.


One preferred embodiment concerns DPPs of general formula III wherein residues A^1 and A^2
15 are different from phenyl.

The DPPs of the general formula III show a high heat stability, a good solubility in polymers, hydrocarbon based fuels, lubricants, and water, a high light stability, and the ability to be used in plastics, especially polyamides, without decomposition and loss of lightfastness, and
20 in paints; and can show photo- and electroluminescence as well as solid state fluorescence. The residues A^1 and A^2 are in general selected from C_1 - C_{18} alkyl, C_2 - C_{18} alkenyl, C_2 - C_{18} alkynyl, C_5 - C_8 cycloalkyl, such as cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, in particular cyclohexyl, C_5 - C_8 cycloalkenyl, such as cyclopentenyl, cyclopentadienyl and cyclohexenyl, in particular cyclohex-3-enyl, aryl and heteroaryl.


25 Diketopyrrolopyrroles, wherein A^1 and A^2 are radicals of the formula



9




R^1 and R^2 are independently of each other hydrogen, halogen, C_1 - C_{18} alkyl, C_1 - C_{18} alkoxy, C_1 - C_{18} alkylmercapto, di(C_1 - C_{18} alkyl)amino, C_1 - C_{18} alkylamino, C_1 - C_{18} alkoxycarbonyl, C_1 - C_{18} alkylaminocarbonyl, -CN, -NO₂, trifluoromethyl, C_5 - C_8 cycloalkyl, -C=N-(C_1 - C_{18} alkyl),



piperazinyl, pyrrolyl, oxazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, morpholinyl, piperidinyl or pyrrolidinyl, -CONX⁵X⁶, -C(O)OX⁷ or -SO₂X⁹; wherein X⁵ and X⁶ are hydrogen, linear or branched C_{1-10} -alkyl, C_{5-10} -cycloalkyl or C_{6-10} -aryl, X⁷ is hydrogen, linear or branched C_{1-10} -alkyl, C_{5-10} -cycloalkyl or C_{6-10} -aryl, X⁹ is hydrogen, linear or branched C_{1-10} -alkyl, C_{5-10} -cycloalkyl, C_{7-10} -aralkyl, C_{6-10} -aryl or -NX¹⁰X¹¹, wherein X¹⁰ and X¹¹ are hydrogen, linear or branched C_{1-10} -alkyl, C_{7-10} -aralkyl or C_{6-10} -aryl,

10 G is -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH=N-, -N=N-, -O-, -S-, -SO-, -SO₂-, -CONH- or -NR⁷-, R^3 and R^4 are independently of each other hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_{18} alkoxy or -CN, R^5 and R^6 are independently of each other hydrogen, halogen or C_1 - C_6 alkyl, and R^7 is

15 hydrogen or C_1 - C_6 alkyl are preferred, wherein radicals of the formula




wherein R^1 and R^2 are independently of each other hydrogen, chloro, bromo, C_1 - C_4 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylamino, phenyl or CN,

20 G is -O-, -NR⁷-, -N=N- or -SO₂-, R^3 and R^4 are hydrogen, and


10

R^7 is hydrogen, methyl or ethyl are further preferred and diketopyrrolopyrrole analogues, wherein A^1 and A^2 are radicals of the formula




wherein R^1 and R^2 are independently of each other hydrogen, methyl, tert-butyl, chloro, bromo, phenyl or CN are particularly preferred for the preparation of inks, colorants, pigmented plastics for coatings, non-impact-printing material, color filters, cosmetics, polymeric ink particles, toners.

In the case of electroluminescence applications the following residues are preferred for A^1 and A^2 :




15


11




- wherein R^{21} , R^{22} , R^{23} , R^{25} and R^{26} are independently of each other hydrogen, C_1 - C_8 alkyl, a hydroxyl group, a mercapto group, C_1 - C_8 alkoxy, C_1 - C_8 alkylthio, halogen, halo- C_1 - C_8 alkyl, a cyano group, an aldehyde group, a ketone group, a carboxyl group, an ester group, a carbamoyl group, an amino group, a nitro group, a silyl group or a siloxanyl group and R^{24} is a C_1 - C_8 alkyl group. Preferably R^{21} , R^{22} , R^{23} , R^{25} and R^{26} are independently of each other hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkoxy or C_1 - C_8 alkylthio, wherein the following residues are particularly preferred:

10






- The residue A^3 is in general selected from hydrogen, C_1 - C_{18} alkyl, cyanomethyl, Ar^3 ,
 5 $-CR^{30}R^{31}-(CH_2)_m-Ar^3$ or $Y-R^{32}$, wherein R^{30} and R^{31} independently of each other stand for
 hydrogen or C_1 - C_4 alkyl, or phenyl which can be substituted up to three times with C_1 - C_3 alkyl,
 Ar^3 stands for aryl, in particular phenyl or 1- or 2-naphthyl, C_5 - C_8 cycloalkyl, such as
 cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, in particular cyclohexyl, C_5 -
 C_8 cycloalkenyl, in particular cyclopentenyl, cyclopentadienyl and cyclohexenyl, or heteroaryl,
 10 which can be substituted one to three times with C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen or phenyl,
 which can be substituted with C_1 - C_8 alkyl or C_1 - C_8 alkoxy one to three times, and m stands for
 0, 1, 2, 3 or 4, Y is $-C(O)-$, $-C(O)O-$, $-C(O)NH-$, $-SO_2NH-$ or $-SO_2-$ and R^{32} is C_1 - C_{18} alkyl, Ar^3 ,
 or aralkyl.
- 15 A^3 is preferably hydrogen, C_1 - C_8 alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-
 butyl, isobutyl, tert.-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2,2-dimethylpropyl, n-hexyl, n-heptyl, n-
 octyl, 1,1,3,3-tetramethylbutyl and 2-ethylhexyl, $Y-R^{32}$ wherein Y is $-C(O)-$ and R^{32} is




$-(CH_2)_m-Ar$ wherein m is 1 and Ar is a group of the formula

13




which can be substituted one to three times with C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen or phenyl. Examples of preferred residues Ar are



- 5 wherein R^{50} and R^{51} are independently of each other methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-butyl, isobutyl, tert.-butyl, methoxy, ethoxy, isopropoxy, tert.-butoxy or chlorine.

The residue A^4 is in general selected from C_1 - C_{18} alkyl or Ar^3 , in particular Ar^3 , wherein A^4 is preferably



10 which can be substituted one to three times with C_1 - C_8 alkyl, C_1 - C_8 alkoxy, halogen or phenyl.


- The fuopyrroles of the formula I are intermediates in the process for the preparation of the diketopyrrolopyrroles of the formula III and, as described in WO03022848, can be used as crystal growth regulators, wherein the term "regulating the crystal growth" refers to controlling the synthesis of pigment particles to have a suitable pigmentary size and/or a narrow particle size distribution as well as directing the growth of the crystals to generate particles of a specifically desired shape, such as platelet, needle, cubic, leaflet, prismatic and other geometric forms and/or of a specifically desired rheology. Consequently, the better control of the crystal growth affords samples with a narrower particle size distribution and/or a better crystal shape, or both. The effect can be influenced by the chemical structure of the organic pigment, the selection of the reaction media and the concentration and chemical structure of the inventive particle growth regulator.

- 25 If used as crystal growth regulator the fuopyrroles of the formula I are present in amount of from about 0.1-20%, especially from 1.0 to 10.0%, based on primary pigment weight. Although DPPs are preferred as primary pigment, the use of diverse pigment moieties is likewise available where the respective pigments are color compatible.


Examples of applicable organic primary pigments are: anthraquinone, phthalocyanine, perinone, perylene, dioxazine, diketopyrrolopyrrole, thioindigo, isoindoline, isoindolinone, quinacridone, quinacridonequinone, flavanthrone, indanthrone, anthrapyrimidine or quinophthalone pigments, and solid solutions comprising these pigments. Preferred organic pigments are quinacridones, phthalocyanines, anthraquinones, perylenes, diketopyrrolopyrroles, isoindolinones and indanthrones.

When the pigment compositions are prepared, the diketopyrrolopyrrole analogues of the formula I can be added during the pigment synthesis, during the fine dispersion process, before or after a finishing process by methods well-known in the art (cf. WO03022848).

Furopyrroles of the formula I, wherein A^1 and A^2 are radicals of the formula




R^1 and R^2 are independently of each other hydrogen, halogen, C_1 - C_{18} alkyl, C_1 - C_{18} alkoxy, C_1 - C_{18} alkylmercapto, C_1 - C_{18} alkylamino, C_1 - C_{18} alkoxycarbonyl, C_1 - C_{18} alkylaminocarbonyl, -CN, -NO₂, trifluoromethyl, C_5 - C_8 cycloalkyl, -C=N-



piperazinyl, pyrrolyl, oxazolyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, morpholinyl, piperidinyl or pyrrolidinyl, -CONX⁵X⁶, -C(O)OX⁷, -SX⁸, -SOX⁸, or -SO₂X⁸; wherein X⁵ and X⁶ are hydrogen, linear or branched C_{1-10} -alkyl, C_{5-10} -cycloalkyl or C_{6-10} -aryl, X⁷ is hydrogen, linear or branched C_{1-10} -alkyl, C_{5-10} -cycloalkyl or C_{6-10} -aryl, X⁸ is hydrogen, linear or branched C_{1-18} -alkyl, C_{5-10} -cycloalkyl, C_{7-10} -aralkyl, C_{6-10} -aryl or -NX¹⁰X¹¹, wherein X¹⁰ and X¹¹ are hydrogen, linear or branched C_{1-10} -alkyl, C_{7-10} -aralkyl or C_{6-10} -aryl,

15

G is $-\text{CH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{CH}=\text{N}-$, $-\text{N}=\text{N}-$, $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{CONH}-$ or $-\text{NR}^7-$, R^3 and R^4 are independently of each other hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_{18} alkoxy or $-\text{CN}$, R^5 and R^6 are independently of each other hydrogen, halogen or C_1 - C_6 alkyl, and R^7 is hydrogen or C_1 - C_6 alkyl are preferred, wherein radicals of the formula




wherein X^5 and X^6 are hydrogen, linear or branched C_{1-4} -alkyl, X^9 is hydrogen, linear or branched C_{1-18} -alkyl, C_{7-10} -aralkyl, C_{6-10} -aryl or $-\text{NX}^{10}\text{X}^{11}$, wherein X^{10} and X^{11} are hydrogen, linear or branched C_{1-10} -alkyl, C_{7-10} -aralkyl or C_{6-10} -aryl;

G is $-\text{O}-$, $-\text{NR}^7-$, $-\text{N}=\text{N}-$, $-\text{S}-$, $-\text{SO}-$ or $-\text{SO}_2-$,

R^3 and R^4 are hydrogen, and

R^7 is hydrogen, methyl or ethyl are further preferred and diketopyrrolopyrrole analogues,


15 wherein A^1 and A^2 are radicals of the formula




wherein R^1 and R^2 are independently of each other hydrogen, C_{1-4} -alkyl, such as methyl or tert-butyl, halogen, such as chloro or bromo, C_{1-4} -alkoxy or C_{1-4} -thioalkyl, phenyl or CN or $-\text{SO}_2\text{X}^9$, wherein X^9 is C_{1-4} -alkyl, phenyl, benzyl or $\text{NX}^{10}\text{X}^{11}$, wherein X^{10} and X^{11} are

20 hydrogen, C_{1-4} -alkyl, benzyl or phenyl are particularly preferred.

A^3 is preferably hydrogen, C_1 - C_8 alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-butyl, isobutyl, tert.-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2,2-dimethylpropyl, n-hexyl, n-heptyl, n-octyl, 1,1,3,3-tetramethylbutyl and 2-ethylhexyl, Y-R^{32} wherein Y is $-\text{C}(\text{O})-$ and R^{32} is



$\text{-(CH}_2\text{)}_m\text{-Ar}$ wherein m is 1 and Ar is a group of the formula



which can be substituted one to three times with $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_1\text{-C}_8$ alkoxy, halogen or phenyl.

5

$\text{C}_1\text{-C}_{18}$ alkyl is typically linear or branched - where possible - and examples of $\text{C}_1\text{-C}_{18}$ alkyl are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-butyl, isobutyl, tert.-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2,2-dimethylpropyl, n-hexyl, n-heptyl, n-octyl, 1,1,3,3-tetramethylbutyl and 2-ethylhexyl, n-nonyl, decyl, undecyl, dodecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl and octadecyl. $\text{C}_1\text{-C}_8$ alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-butyl, isobutyl, tert.-butyl, n-pentyl, 2-pentyl, 3-pentyl, 2,2-dimethylpropyl, n-hexyl, n-heptyl, n-octyl, 1,1,3,3-tetramethylbutyl and 2-ethylhexyl is preferred. $\text{C}_1\text{-C}_4$ alkyl such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.-butyl, isobutyl or tert.-butyl is particularly preferred. The term "C₂-C₁₈alkenyl group" means an unsaturated linear or branched aliphatic hydrocarbon group containing one or more double bonds, in particular C₂₋₈-alkenyl, such as vinyl, allyl, 2-propen-2-yl, 2-buten-1-yl, 3-buten-1-yl, 1,3-butadien-2-yl, 2-penten-1-yl, 3-penten-2-yl, 2-methyl-1-buten-3-yl, 2-methyl-3-buten-2-yl, 3-methyl-2-buten-1-yl and 1,4-pentadien-3-yl. The term "C₂-C₁₈alkynyl group" means an unsaturated aliphatic hydrocarbon group containing a triple bond, in particular C₂-C₈-alkynyl such as ethynyl, 1-propyn-1-yl, 2-butyne-1-yl, 3-butyne-1-yl, 2-pentyne-1-yl and 3-pentyne-2-yl.

20

Examples of $\text{C}_1\text{-C}_{18}$ alkoxy, which can be linear or branched, are methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec.-butoxy, isobutoxy, tert.-butoxy, n-pentoxy, 2-pentoxy, 3-pentoxy, 2,2-dimethylpropoxy, n-hexoxy, n-heptoxy, n-octoxy, 1,1,3,3-tetramethylbutoxy and 2-ethylhexoxy, wherein $\text{C}_1\text{-C}_4$ alkoxy such as methoxy, ethoxy, n-propoxy, isopropoxy,

25

n-butoxy, sec.-butoxy, isobutoxy and tert.-butoxy is preferred. Examples of

$\text{C}_1\text{-C}_{18}$ alkylmercapto are the same groups as mentioned for the alkoxy groups, except that the oxygen atom of the ether linkage is replaced by a sulphur atom. Examples and preferences for $\text{C}_1\text{-C}_{18}$ alkyl in $\text{C}_1\text{-C}_{18}$ alkylamino and $\text{C}_1\text{-C}_{18}$ alkylaminocarbonyl are the same as mentioned for $\text{C}_1\text{-C}_{18}$ alkyl. Examples and preferences for $\text{C}_1\text{-C}_{18}$ alkoxy in

30

$\text{C}_1\text{-C}_{18}$ alkoxycarbonyl are the same as mentioned for $\text{C}_1\text{-C}_{18}$ alkoxy.

The term "aryl group" is typically $\text{C}_6\text{-C}_{24}$ aryl, such as phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, phenanthryl, terphenyl, pyrenyl, 2- or 9-fluorenyl or anthracenyl, preferably $\text{C}_6\text{-}$

C₁₂aryl such as phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, which may be unsubstituted or substituted.

The term "aralkyl group" is typically C₇-C₂₄aralkyl, such as benzyl, 2-benzyl-2-propyl, β-phenylethyl, α,α-dimethylbenzyl, ω-phenylbutyl, ω,ω-dimethyl-ω-phenylbutyl,

5 ω-phenyldodecyl, ω-phenyloctadecyl, ω-phenyleicosyl or ω-phenyldocosyl, preferably C₇-C₁₈aralkyl such as benzyl, 2-benzyl-2-propyl, β-phenylethyl, α,α-dimethylbenzyl, ω-phenylbutyl, ω,ω-dimethyl-ω-phenylbutyl, ω-phenyldodecyl or ω-phenyloctadecyl, and particularly preferred C₇-C₁₂aralkyl such as benzyl, 2-benzyl-2-propyl, β-phenyl-ethyl, α,α-dimethylbenzyl, ω-phenyl-butyl, or ω,ω-dimethyl-ω-phenyl-butyl, in which both the
10 aliphatic hydrocarbon group and aromatic hydrocarbon group may be unsubstituted or substituted.

Examples of C₅-C₈cycloalkyl are cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, which may be unsubstituted or substituted. The term "C₅-C₈cycloalkenyl group" means an unsaturated alicyclic hydrocarbon group containing one or more double bonds, such as
15 cyclopentenyl, cyclopentadienyl and cyclohexenyl, which may be unsubstituted or substituted.

The term "heteroaryl" is a ring with five to seven ring atoms, wherein nitrogen, oxygen or sulfur are the possible hetero atoms, and is typically an unsaturated heterocyclic radical with five to 18 atoms having at least six conjugated π-electrons such as thienyl, benzo[b]thienyl,
20 dibenzo[b,d]thienyl, thianthrenyl, furyl, furfuryl, 2H-pyranyl, benzofuranyl, isobenzofuranyl, dibenzofuranyl, phenoxythienyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, bipyridyl, triazinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolizynyl, isoindolyl, indolyl, indazolyl, purinyl, quinolizynyl, quinolyl, isoquinolyl, phthalazinyl, naphthyridinyl, quinoxalynyl, quinazolinyl, cinnolynyl, pteridinyl, carbazolyl, carbolinyl, benzotriazolyl, benzoxazolyl, phenanthridinyl, acridinyl,
25 perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl or phenoxazinyl.

Examples of a halogen atom are fluorine, chlorine, bromine and iodine.

If the above-mentioned substituents can be substituted, possible substituents are C₁-C₈alkyl, a hydroxyl group, a mercapto group, C₁-C₈alkoxy, C₁-C₈alkylthio, halogen, halo-C₁-C₈alkyl, a
30 cyano group, an aldehyde group, a ketone group, a carboxyl group, an ester group, a carbamoyl group, an amino group, a nitro group, a silyl group or a siloxanyl group.

As described in WO03022848 the DPPs of the general formula III can be used for the preparation of
35 inks for printing inks in printing processes, for flexographic printing, screen printing, packaging printing, security ink printing, intaglio printing or offset printing, for pre-press

stages and for textile printing, for office, home applications or graphics applications, such as for paper goods, for example, for ballpoint pens, felt tips, fiber tips, card, wood, (wood) stains, metal, inking pads or inks for impact printing processes (with impact-pressure ink ribbons), for the preparation of


- 5 colorants for coating materials, for industrial or commercial use, for textile decoration and industrial marking, for roller coatings or powder coatings or for automotive finishes, for high-solids (low-solvent), water-containing or metallic coating materials or for pigmented formulations for aqueous paints, for the preparation of
- 10 pigmented plastics for coatings, fibers, platters or mold carriers, for the preparation of non-impact-printing material for digital printing, for the thermal wax transfer printing process, the ink jet printing process or for the thermal transfer printing process, and also for the preparation of
- color filters, especially for visible light in the range from 400 to 700 nm, for liquid-crystal displays (LCDs) or charge combined devices (CCDs) or for the preparation of
- 15 cosmetics or for the preparation of polymeric ink particles, toners, dye lasers, dry copy toners liquid copy toners, or electrophotographic toners, and electroluminescent devices.

- 20 The following examples illustrate various embodiments of the invention, but the scope of the invention is not limited thereto.

- The microwave generator used was a CEM Discover® model, with a circular single mode cavity design, that focuses the microwave radiation on the sample. The sample was contained in a sealed glass tube, whereby the pressure was allowed to increase to a
- 25 maximum of $20.69 \cdot 10^5$ Pa (300 p.s.i.). The maximum operating power of this device was 300 watts. ^1H and ^{13}C NMR spectra were obtained at 300 and 75 MHz respectively, and coupling constants are in Hz. Mass spectral measurements were obtained using chemical ionisation at 70 eV, with isobutane as carrier gas.

Examples

Example 1

5 **3,6-Diphenylfuro[3,4-c]pyrrole-1,4-dione (2)**

Ethyl 4-benzoyl-4,5-dihydro-5-oxo-2-phenylpyrrole-3-carboxylate **1** (99.5 mg, 0.296 mmol, prepared as previously reported in WO03022848) was irradiated with microwave radiation (at a frequency of 2 to 45 GHz, and a forward power of 300 Watts) without solvent, heating to 250 °C for 10 minutes. The crude product was then allowed to cool, methanol was added and the solid filtered off and washed with methanol. This gave the furopyrrole **2** as an orange solid (73 mg, 86 %). Decomp > 300 °C.

δ_H (DMSO d_6) 11.87 (1H, s, NH), 8.12 – 8.23 (4H, dm, Ar-H) and 7.48 – 7.54 (6H, m, Ar-H);


δ_C (DMSO d_6) 161.4, 159.3 (2 x C=O), 152.2, 148.1 (2 x quat.), 132.8, 132.6, 129.1 (2C), 128.0, 127.0 (6 x Ar C-H), 126.8, 126.4, 115.8, 102.8 (4 x quat.).

15

Comparative Example 1 (Example 1 of WO03022848)

A mixture of ethyl 4-benzoyl-4,5-dihydro-5-oxo-2-phenylpyrrole-3-carboxylate **1** (10 g, 0.0299 mol) and Dowtherm A (200 ml) was heated to 230-240 °C under nitrogen for 64 h. The solution was then cooled to 25 °C and added dropwise to petrol ether 40-60 (300 ml) upon which a fluorescent orange solid precipitated. This was filtered off, washed with further hexane and dried *in vacuo*. Yield 3.48 g (40 %).

20

Example 2**5 a) *p*-Bromobenzoyl Chloride (3)**

p-Bromobenzoic acid was purified by dissolving in NaOH (aq) and washing the solution with dichloromethane, followed by acidification of the aqueous layer with dilute aqueous HCl, and extraction with EtOAc. The acid (4.00 g, 0.0182 mol), oxalyl chloride (4.634 g, 3.185 ml, 0.0364 mol), and a catalytic amount of DMF was stirred overnight at room temperature in DCM (40 ml). Evaporation of the solvents and excess reagents gave the acid chloride 3 as an off-white solid. m.p. 38–40 °C.

b) Ethyl 4-(*p*-bromobenzoyl)-4,5-dihydro-5-oxo-2-phenylpyrrole-3-carboxylate (5)

To sodium hydride (590 mg, 14.75 mmol) was added THF (40 ml), and the pyrrolinone ester 4 (852 mg, 3.69 mmol). After stirring for 30 mins at room temperature, a solution of *p*-bromobenzoyl chloride (809.5 mg, 3.69 mmol) in THF (10 ml) and a catalytic amount of DMAP, was added and the mixture was stirred at room temperature overnight. 10 % HCl (aq) was added, and the organic component extracted with diethyl ether. Concentration *in vacuo* and recrystallisation from ethanol gave the enol 5 as a yellow crystalline solid (665 mg, 44 %). M.p. 189 °C; δ_{H} (DMSO d_6) 11.90 (1H, s, NH), 7.72 – 7.82 (2H, m, ArH), 7.58 – 7.66 (4H, m, ArH), 7.42 – 7.53 (3 H, m, ArH), 3.72 (2H, q, CH₂CH₃) and 0.9 (3H, t, CH₂CH₃); m/z 416

(M+1 ⁸¹Br, 100 %), 414 (M+1 ⁷⁹Br, 96 %) 347, 319, 317, 296

c) 3-(*p*-Bromophenyl)-6-phenyl furo[3,4-*c*]pyrrole-1,4-dione (6)


The *p*-bromobenzoylpyrrolinone ester 5 (154 mg, 0.37 mmol) was irradiated with microwave radiation (at a frequency of 2 to 45 GHz, and a forward power of 300 Watts) without solvent, heating to 250 °C for 10 minutes. The crude product was then allowed to cool, methanol was added and the solid filtered off and washed with methanol. This gave the furopyrrole 6 as a red solid (129 mg, 94 %). M.p. 295 °C (subl., decomp.); δ_{H} (DMSO *d*₆) 11.88 (1H, s, NH), 8.13 – 8.17 (2H, m, Ar-*H*), 7.98 and 7.66 (2 x 2H, AA'BB', *J* 8.7, C₆H₄) and 7.43 – 7.47 (3H, m, Ar-*H*); *m/z* 370 (M+1 ⁸¹Br, 94 %) and 368 (M+1 ⁷⁹Br, 100 %).

d) 5-Methyl-3-(*p*-bromophenyl)-6-phenylfuro[3,4-*c*]pyrrole-1,4-dione (7)

A mixture of furopyrrole 6 (1.5 g, 4.08 mmol), methyl tosylate (1.14 g, 6.12 mmol), potassium carbonate (1.13 g, 8.16 mmol) and dimethylformamide was stirred at room temperature overnight. Water was then added, and the organic component extracted with DCM. The solvent was removed, and washing with water then methanol gave the methylated compound 7 as a red solid (0.831 g, 53 %), m.p. 215-216 °C. δ_{H} (CDCl₃) 8.19 and 7.61 (each 2H, AA'BB', *p*-C₆H₄Br), 7.78-7.73 (2H, m, *o*-Ph), 7.54-7.50 (3H, m, *m/p*-Ph) and 3.38 (3H, s, NCH₃). λ_{max} (abs) (DCM)/nm 454 (ϵ 15,878)

e) 2-Methyl-5-phenyl-6-(*p*-Bromophenyl)-3-phenylpyrrolo[3,4-*c*]pyrrole-1,4-dione (8)

A mixture of furopyrrole 7 (300 mg, 0.79 mmol), aniline (146 mg, 1.57 mmol), DCC (323 mg, 1.57 mmol), trifluoroacetic acid (2-3 drops) and DCM was stirred at room temperature for 144 hours. The solvent was removed, and washing with methanol gave the pyrrolopyrrole 8 as a red solid (173 mg, 55 %), m.p. 255-256 °C. δ_{H} (CDCl₃) 7.88-7.83 (2H, m, Ar-*H*), 7.49-7.43 (5H, m, Ar-*H*), 7.40-7.26 (5H, m, Ar-*H*), 7.12-7.07 (2H, m, Ar-*H*) and 3.35 (3H, s, NCH₃)

Example 3**a) Ethyl 4-(p-nitrobenzoyl)-4,5-dihydro-5-oxo-2-phenylpyrrole-3-carboxylate (11)**

- 5 The pyrrolinone ester **9** (6.35 g, 27.5 mmol) was added to a mixture of sodium hydride (2.0 g, 82.5 mmol) and THF (1 litre), and this was stirred at room temperature for 15 minutes. *p*-Nitrobenzoyl chloride **10** was then added, and the mixture was stirred overnight. Methanol was added, followed by water and the mixture acidified with HCl. The organic component was extracted with diethyl ether and the solvent evaporated. Washing with methanol gave the nitro compound **11** as a yellow solid (6.31 g, 60 %). δ_{H} (DMSO- d_6) 11.95 (1H, s, NH), 8.30 and 7.84 (each 2H, AA'BB', Ar), 7.56-7.50 (2H, m, *o*-Ph), 7.45-7.35 (3H, m, *m/p*-Ph), 3.62 (2H, q, OCH_2CH_3) and 0.75 (3H, t, OCH_2CH_3). m/z (ESI -ve) 380 (22 %, M^+), 379 [100 %, $(\text{M} - 1)]^+$

15 b) 3-(p-Nitrophenyl)-6-phenylfuro[3,4-c]pyrrole-1,4-dione (12)

- The *p*-nitrobenzoylpyrrolinone ester **11** (300 mg, 0.90 mmol) was irradiated with microwave radiation without solvent, heating to 270 °C for 15 minutes. The crude product was then allowed to cool, methanol was added and the solid filtered off and washed with methanol. This gave the *furo*pyrrole **12** as a red solid (230 mg, 87 %).
- 20 δ_{H} (DMSO- d_6) 12.15 (1H, s, NH), 8.38 (4H, s, $p\text{-C}_6\text{H}_4\text{NO}_2$), 8.34 – 8.28 (2H, m, *o*-Ph) and 7.69 – 7.58 (3H, m, *m/p*-Ph). m/z (ESI -ve) 334 (21 %, M^+), 333 [100 %, $(\text{M} - 1)]^+$

c) 5-Methyl-3-(*p*-nitrophenyl)-6-phenylfuro[3,4-*c*]pyrrole-1,4-dione (13)

A mixture of furopyrrole **12** (0.9 g, 2.7 mmol), methyl tosylate (750 mg, 4.04 mmol), potassium carbonate (1 g, 7.2 mmol) and dimethyl formamide was stirred at room temperature overnight. Water was then added, and the organic component extracted with DCM. The solvent was removed, and washing with water then methanol gave the methylated compound **13** as a red solid (0.652 g, 70 %), m.p. 253-255 °C. δ_{H} (CDCl₃) 8.55 and 8.38 (4H, AA'BB', Ar), 7.88-7.84 (2H, m, *o*-Ph), 7.65-7.60 (3H, m, *m/p*-Ph) and 3.49 (3H, s, NCH₃). λ_{max} (abs) (DCM)/nm 482 (ϵ 17,462)


10

d) 2-Methyl-5'-phenyl-6-(*p*-nitrophenyl)-3-phenylpyrrolo[3,4-*c*]pyrrole-1,4-dione (14)

A mixture of furopyrrole **13** (100 mg, 0.29 mmol), aniline (53 mg, 0.57 mmol), DCC (118 mg, 0.57 mmol), trifluoroacetic acid (2-3 drops) and DCM was stirred at room temperature for 72 hours. The solvent was removed, and washing with methanol gave the pyrrolopyrrole **14** as a red solid (63 mg, 52 %), m.p. 233-235 °C. δ_{H} (CDCl₃) 8.16 and 7.81 (each 2H, AA'BB', *p*-C₆H₄NO₂), 7.98-7.93 (2H, m, *o*-Ph), 7.60-7.55 (3H, m, *m/p*-Ph), 7.44-7.36 (3H, m, *m/p*-Ph), 7.20-7.15 (2H, m, *o*-Ph) and 3.45 (3H, s, NCH₃). λ_{max} (abs) (DCM)/nm 493 (ϵ 14,014)

15

24

Example 4**a) Ethyl 4,5-dihydro-5-oxo-1,2-diphenylpyrrole-3-carboxylate (16)**

- 5 Aniline (2.65 g, 2.59 ml, 0.0285 mmol) was added to a solution of ethyl 5-oxo-2-phenyl-4,5-dihydro-furan-3-carboxylate **15** (made via the literature method described in F. Gaudemar-Bardone, M. Mladenova, R. Couffignal, *Synthesis*, 1985, 1043) (6.0 g, 0.0259 mmol) and acetic acid (100 ml), and the solution heated to reflux for 3 hours. The solution was then cooled, diluted with water and extracted with diethyl ether. The organic extracts were dried and concentrated. Column chromatography (silica gel, eluent dichloromethane) gave the lactam **16** as a colourless solid (6.9 g, 87 %), m.p. 129-130 °C. δ_H (CDCl₃) 7.32-7.15 (8H, m, Ar-H), 6.98-6.93 (2H, m, o-Ph-N), 4.08 (2H, q, OCH₂CH₃, J 6.9), 3.67 (2H, s, CH₂) and 1.11 (3H, t, OCH₂CH₃, J 6.9)
- 10

b) Ethyl 4-benzoyl-4,5-dihydro-5-oxo-1,2-diphenylpyrrole-3-carboxylate (17)

A solution of pyrrolinone ester **16** (1.76 g, 5.74 mmol) in tetrahydrofuran (100 ml) was cooled

to -78°C , and a 1.0 M solution of lithium hexamethyldisilazide (17.2 ml, 17.2 mmol) in THF was added. After 5 minutes, benzoyl chloride (0.97 g, 0.79 ml, 6.89 mmol) was added, and the solution stirred for 30 mins. Methanol was added, and the solution warmed to room temperature. The mixture was acidified (aqueous HCl) and extracted with diethyl ether. The ether extracts were dried and concentrated. Column chromatography (silica gel, eluent dichloromethane) gave the enol **17** as a yellow solid (1.74 g, 74 %), m.p. $137-139^{\circ}\text{C}$. δ_{H} (CDCl_3) 7.75-7.68 (2H, m, Ar), 7.54-7.44 (3H, m, Ar), 7.34-7.19 (8H, m, Ar), 7.14-7.07 (2H, m, Ar), 3.54 (2H, q, CH_2 , J 7.2) and 0.65 (3H, t, CH_3 , J 7.2)

c) 3,5,6-triphenyl-1H-furo[3,4-c]pyrrole-1,4(5H)-dione (18)

Benzoyl pyrrolinone ester **17** (74 mg) was irradiated with microwave radiation (at 300 Watts) without solvent, heating to 200°C for 10 minutes. The crude product was then allowed to cool, methanol was added and the solid filtered off and washed with methanol. This gave the furopyrrole **18** as an orange solid (34 mg, 52 %), m.p. $230-232^{\circ}\text{C}$ (lit. [H. Langhals, T. Grundei, T. Potrawa, K. Polborn, *Liebigs Ann. Chem.*, 1996, 679] $230-232^{\circ}\text{C}$). δ_{H} (CDCl_3) 8.48-8.42 (2H, m, Ar-H) and 7.61-7.20 (13H, m, Ar-H)

d) 2,3,5,6-Tetraphenyl-2,5-dihydropyrrolo[3,4-c]pyrrole-1,4-dione (19)

Can be prepared starting from the intermediate **18** as described in H. Langhals, T. Grundei, T. Potrawa, K. Polborn, *Liebigs Ann. Chem.*, 1996, 679.

e) 2,3,6-triphenylpyrrolo[3,4-c]pyrrole-1,4-dione (21)

N-phenyl pyrrolinone ester **16** (663 mg, 2.16 mmol) and benzonitrile (446 mg, 440 μl , 4.3 mmol) were added successively to a solution of sodium t-amyl oxide [from sodium (150 mg, 6.5 mmol) and t-amyl alcohol (4.0 ml)], and the mixture heated to reflux for 6 hours. The mixture was then cooled, and acidified (dilute aqueous HCl) and extracted with dichloromethane. The organic extracts were then dried and the solvent evaporated. Precipitation from methanol, followed by filtration gave the triphenyl pyrrolopyrrole **21** as a bright orange solid (18 mg, 3 %), m.p. 390°C (decomp). δ_{H} ($\text{DMSO}-d_6$) 11.54 (1H, s, NH), 8.55 (2H, m, Ar-H), 7.57-7.67 (5H, m, Ar-H), 7.38-7.54 (6H, m, Ar-H), 7.29-7.33 (2H, m, Ar-H); m/z (LCMS) 363.96 (30%, M), 362.95 (100 %, M-H); λ_{max} abs (DMSO)/nm 269 (ϵ /24,550), 303 (15,720) 470 (22,580) and 498 (23,640)